

Doxitec

FOR VETERINARY USE

Action: Antibiotic for dogs and cats.

Presentation:

Carton box with 2 blisters containing 8 tablets of 50 mg each
Carton box with 2 blisters containing 8 tablets of 100 mg each
Carton box with 2 blisters containing 8 tablets of 200 mg each

Formula:

Each tablet contains:

	50mg	100mg	200mg
Doxycycline hyclate	56.5 mg	113.0 mg	226.0 mg
(Equivalent to 50 mg / 100 mg / 200 mg of doxycycline)			
Excipients q.s.	150.0 mg	300.0 mg	600.0 mg

Doxitec is a doxycycline-based antibiotic, a modern and effective antimicrobial belonging to tetracyclines group. **Doxitec** is indicated for treatment of a variety of extracellular and intracellular infections in dogs and cats, especially canine Ehrlichiosis. **Doxitec** acts on respiratory, urinary and gastrointestinal infections caused by *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus pyogenes*, *Klebsiella pneumoniae*, *Salmonella typhimurium*, *Shigella flexnerie*, *Bordetella bronchiseptica*.

Dosage:

For Dogs:

Treatment of infections:

Doxitec is administered by oral route, at a dosage of 5 mg/kg body weight, i.e.:

- 1 tablet of 50 mg each 10 kg body weight.
- 1 tablet of 100 mg each 20 kg body weight.
- 1 tablet of 200 mg each 40 kg body weight.

Dogs weighing over 50 kg should be treated following the proportion as above described.

This dose should be administered each 12 hours for 7 consecutive days or at the discretion of Veterinarian. **Doxitec** can be administered with feed or dissolved in liquid.

Treatment of Canine Ehrlichiosis:

Dose: 10 mg/kg.

This dose should be administered each 12 hours for 21 consecutive days or at the discretion of Veterinarian.

Maintaining treatment for 48 hours after the remission of symptoms is recommended.

Do not administer product for pregnant females during the last third of gestation, as so as for young animals during dentition development.

For Cats:

Doxitec is administered by oral route, at the dose of 10 mg/kg body weight, ie:

- 1 tablet of 50 mg each 5 kg body weight.

- 1 tablet of 100 mg each 10 kg body weight.

Cats weighing over 10 kg should be treated following the proportion as above described. This dose should be administered each 24 hours for 7 consecutive days or at the discretion of Veterinarian, **Doxitec** can be administered with feed or dissolved in liquid.

Maintaining treatment for 48 hours after the remission of symptoms is recommended.

Adverse effects: The main side effects reported were: nausea, vomit, anorexia and/or diarrhea. These effects can be relieved when administering the drug with feed, what does not affect its absorption. These effects are usually mild and transient.

Contraindications: **DOXITEC** is contraindicated for pregnant females during the final third of pregnancy, in lactating dogs or cats and in young animals during dentition development, and in case of known hypersensitivity to tetracyclines.

Precautions: Avoid simultaneous administration of **DOXITEC** with antacids and compounds containing iron and calcium. In evidence of erythema, discontinue use. Avoid human consumption.

Do not reuse packaging. Keep the product in its original packaging.

Store the product at room temperature in a cool, dry place and away from direct sunlight;

KEEP THE PRODUCT OUT OF REACH OF CHILDREN AND PETS.

SOLD UNDER PRESCRIPTION AND APPLICATION UNDER VETERINARIAN DIRECTION.

Licensed by the Ministry of Agriculture with No. 9089/2005 in 2005/05/15.

Responsible Technician:

Dra. Vera de Fátima Costa Calixto - CRF / SP 11.427

License Holder and

Manufacturer:

Syntec do Brasil Ltda.
Estrada Rosa Scarpa, 178 -
Votuparim- Santana de
Parnaíba/SP CEP:
06513-010 CNPJ:
02.177.011/00011-77 A
Brazilian Company

CUSTOMER CARE

SERVICE +55 (11)

4702-5425

sac@syntecvet.com.br

www.syntecvet.com.br

